

## Relation of Oxidant-Antioxidant Status with Glycemic control in type 2 diabetic patients

\*Baydaa Ahmmad Abed

\*National Diabetes Center Al – Mustansiria University

### Abstract:

In diabetes mellitus, persistence of hyperglycemia was reported to cause increased production of oxidative parameters, and decreased antioxidant status. Therefore, the present study considered to evaluate the effect of glycemic control on oxidant-antioxidant status in type 2 diabetic patients. The study examined 145 type 2 diabetic patients, who attend to the National Diabetic Center (Al-Mustansiyriah University). Patients were divided in two groups, as well glycemic controlled (group1[75 patients]) ( $HbA_{1c} < 7\%$ , and  $FSG < 110$  mg/dl) and poorly controlled (group2 [70 patients]) (had  $HbA_{1c} \geq 7\%$ , and  $FSG \geq 110$ ). All patients were reported in the morning after overnight fast, and underwent physical examination and laboratory tests. Height and weight were noted for Body Mass Index (BMI), it was calculated as weight (Km)/height<sup>2</sup> (m<sup>2</sup>).

Laboratory evaluations consisted of measuring fasting serum glucose (FSG), glycated hemoglobin ( $HbA_{1c}$ ), total cholesterol (TC), triglycerides (TG), malondialdehyde (MDA), erythrocyte-Glutation (GSH), ascorbic acid (AA), uric acid (UA), and cerulopalsmine (CP). The results indicate that fasting serum FSG,  $HbA_{1c}$ , (MDA), (TC), and (TG), were increased significantly ( $P < 0.01$  for FSG,  $HbA_{1c}$ , MDA; and  $P < 0.05$  for TC, and TC) in poorly controlled subjects comparing to well controlled. Instead of that there is significant increase in (CP) ( $P < 0.05$ ) in poorly controlled patients. On the other hand there is significant decreased in (GSH, (AA), (UA) ( $P < 0.01$  for GSH, AA; and  $P < 0.05$  for UA) in poorly controlled compared to well controlled diabetic patients. Correlation analysis showed a significant positive correlation in serum  $HbA_{1c}$ -MDA, GSH-AA ( $P < 0.01$ ; and  $P < 0.05$  respectively) in both group, while there is only positive correlation in  $HbA_{1c}$ -FSG ( $P < 0.01$ ) in well controlled patients, and positive correlation in  $HbA_{1c}$ -TC in poorly controlled patients. On the other hand, there is a significant negative correlation in MDA-GSH, MDA-AA, and MDA-UA in both studying group. The present study concludes the increased risk of oxidative stress manifested by increased plasma MDA, and decreased antioxidant levels in poor glycemic control. The study therefore suggests, the estimation of antioxidants levels with other routine investigations may be useful in reduce of the oxidative stress in diabetics.

**Keywords:** Glycemic control.Type2 diabetes.Oxidative stress.

### الخلاصة:

افادت دراسات عديدة ان ارتفاع السكر بالدم للمرضى المصابين بداء السكر من نوع الثاني يؤدي الى زيادة معلمات الاكسدة وتراجع نسبة مضادات الاكسدة. لذلك اجريت هذه الدراسة لتقييم مدى تأثير سيطرة نسبة السكر في الدم لمرضى النوع الثاني على معلمات الاكسدة ومضادات الاكسدة. شملت هذه الدراسة 145 مريض مصاب بداء السكر من النوع الثاني من المراجعين للمركز الوطني لعلاج وبحوث السكري/ الجامعة المستنصرية حيث تم تقسيم المرضى الى مجموعتين المجموعة الاولى G1 ذات السيطرة الجيدة (75 مريضا)(نسبة السكر بالدم  $> 110$  وخضاب الدم المعسل  $> 7\%$ ) والمجموعة الثانية G2 ذات السيطرة السيئة(70 مريضا)(نسبة السكر بالدم  $\leq 110$  وخضاب الدم المعسل  $\leq 7\%$ ). اخذت جميع العينات من المرضى بعد صيام 12 ساعة وبعد ان تم فحصهم سريريا ومختبريا. تم حساب معامل كتلة الجسم عن طريق اخذ الوزن والطول(الوزن على مربع الطول بالامتر).اجريت للمرضى التحاليل التالية (مستوى السكر بالدم, خضاب الدم المعسل,الكوليسترول,الدهون الثلاثية,(MDA)malondialdehyde,erythrocyte–Glutation(GSH),حامض الاسكوريك,حامض اليوريك و ceruloplasmine. اظهرت النتائج ان هناك زيادة معنوية في مستوى كل من (السكر في الدم,خضاب الدم المعسل,(MDA) malondialdehyde,الكوليسترول والدهون الثلاثية) حيث كانت هناك زيادة معنوية عالية جدا

( $P < 0.01$ ) في كل من مستوى السكر بالدم، خضاب الدم المعسل، (MDA) malondialdehyde وكانت هناك زيادة معنوية ( $P < 0.05$ ) في كل من الكوليسترول والدهون الثلاثية في المجموعة ذات السيطرة السيئة اذا ما قورنت بالمجموعة ذات السيطرة الجيدة بالإضافة الى ذلك كانت هناك زيادة معنوية في ceruloplasmine (CP) ( $P < 0.05$ ) من ناحية اخرى كان هناك انخفاض معنوي في erythrocyte-GSH وحامض الاسكورب A.A، حامض اليوري U.A ( $P < 0.01$ ) في كل من GSH وحامض الاسكورب A.A و  $P < 0.05$  في حامض اليوريك U.A في المجموعة ذات السيطرة السيئة اذا ما قورنت بالمجموعة ذات السيطرة الجيدة واطهرت تحاليل الارتباط وجود علاقة معنوية موجبة بين ال HbA1C-MDA ( $P < 0.01$ )، A.A-GSH ( $P < 0.05$ ) في المجموعتين بينما كانت هناك علاقة معنوية موجبة واحدة فقط في مرضى السيطرة الجيدة بين (سكر الدم - خضاب الدم المعسل) ( $P < 0.01$ )، وعلاقة موجبة واحدة بين الكوليسترول - خضاب الدم المعسل في مرضى السيطرة السيئة. من ناحية اخرى كانت هناك علاقة معنوية سالبة بين MDA-GSH، MDA-AA، MDA-U.A في كلا المجموعتين.

نستنتج من هذه الدراسة ان زيادة خطر اجهاد الاكسدة وانخفاض مستوى مضادات الاكسدة يكون واضحا بزيادة MDA عند مرضى السكري ذوي السيطرة السيئة لمستوى السكر في الدم. من خلال هذه النتائج نقترح الاعتماد على قياس نسبة مضادات الاكسدة كجزء من تحاليل الروتينية التي قد تكون مفيدة في الحد من اجهاد الاكسدة في مرضى السكري.

## Introduction:

The main cause of morbidity and mortality in both type 1 and type 2 diabetes (T1DM, T2DM) is the development of micro- and macrovascular complications. Several studies have shown a direct relation between glucose exposure, as measured by hemoglobin A1c (HbA1c) level, and complications of diabetes<sup>[1]</sup>. The molecular mechanisms that explain this relation between hyper-glycemia and diabetic complications have not been fully elucidated. A suggested common pathway by which hyperglycemia leads to these complications is the formation of reactive oxygen species (ROS) by glucose overload in the mitochondria. This could lead to vascular damage through several molecular mechanisms<sup>[2]</sup>. However, HbA1c only explains around 25% of the variation in risk of developing complications<sup>[3]</sup>, suggesting the contribution of other factors. It has been suggested that glycemic variability contributes to diabetes complications via formation of ROS<sup>[4]</sup>. Shadia *et. al.*<sup>[5]</sup> showed that persistence of hyperglycemia in diabetes mellitus is the cause to increased production of oxidative parameters as well as defects in antioxidant defense system. A strong relationship between glycemic variability and oxidative stress in poorly regulated type 2

diabetes (T2DM) on oral medication has been reported<sup>[6]</sup>. Free radicals are very reactive chemical species, can cause oxidative injury to the living organisms by attacking the macromolecules like lipids, carbohydrates and nucleic acids. Under normal physiological conditions, there is a critical balance between the generation of oxygen free radicals and antioxidant defense systems used by the organisms, to deactivate and protect themselves against free radical toxicity<sup>[7]</sup>. Impairment in the oxidant/antioxidant equilibrium creates a condition known as oxidative stress. It is known to be a component of molecular and cellular tissue damage mechanism in a wide spectrum of human diseases<sup>[8]</sup>. Mechanisms involved in the increased oxidative stress in diabetes include not only oxygen free radical generation due to non-enzymatic glycosylation (placation), autooxidation of glycation products, but also changes in the tissue content and activity of the antioxidant defense systems<sup>[9]</sup>. Oxidative stress as well as defects in antioxidant defense systems are recognized as causative factors for the development of major diabetic complications<sup>[10]</sup>. A variety of natural antioxidants exist to scavenge oxygen free radicals and prevent oxidative damage to biological membranes. One group of these antioxidants is enzymatic (intracellular), which include

super oxide dismutase, glutathione peroxidase and catalase. In addition to enzymatic antioxidants, the major natural antioxidants, most of them derived from natural sources by dietary intake are vitamin A, vitamin C and vitamin E and carotenoids. Also, numerous small molecules are synthesized or produced within the body that has antioxidant capacity (e.g. glutathione and uric acid), and some metal chelator such as ceruloplasmine and transferrine<sup>[11-13]</sup>. Lipids, the substrate under greatest attack by free radicals, have been the most extensively studied and in diabetic patients are of special interest because their hyperlipidemia (excess substrate) is considered to be a significant risk factor for the development of microvascular complications<sup>[14]</sup>. Peroxidation of fatty acids produces aldehydes such as thiobarbituric acid reacting substances (TBARS). The most commonly measured TBARS is malon dialdehyde<sup>[15]</sup>. A number of studies have evaluated oxidative stress and antioxidant status in type 2 diabetic patients<sup>[9,11]</sup>. this study was planned to evaluate serum lipid peroxidation marker, malondialdehyde (an oxidant) and selected antioxidant (erythrocyte-Glutation (GSH), Ascorbic acid (AA), uric acid (UA) and ceruloplasmine (CP) in relation to glycemic control in type 2 diabetic patients.

### **Patients and Methods:**

The current study included 145 previously diagnosed type 2 diabetic patients; their mean age was 47.7±10.68 years, who attend to the National Diabetic Center (Al-Mustansiyriah University). Study cases were divided in two groups according their mean hemoglobin (HbA<sub>1c</sub>), and fasting serum glucose (FSG) values, as good controlled (group1 [75 patients]) (had HbA<sub>1c</sub>< 7%, and FSG < 110 mg/dl) and poorly controlled (group2 [70 patients]) (had HbA<sub>1c</sub>≥ 7%, and FSG ≥ 110) patients with DM, both group were matching with age,

body mass index (BMI). The selected patients with diagnosed cardiovascular disease, kidney disease, liver disease, and those who smoked were excluded from this study; none of the patients were taking vitamin or mineral supplement shortly before or during the study. All patients were reported in the morning after overnight fast, and underwent physical examination and laboratory tests. Height and weight were noted for Body Mass Index (BMI), it was calculated as weight (Kilogram)/height<sup>2</sup> (meter<sup>2</sup>). Laboratory evaluations consisted of measuring fasting serum glucose (FSG), glycated hemoglobin (HbA<sub>1c</sub>), total cholesterol (TC), triglycerides (TG), malon dialdehyde (MDA), erythrocyte-Glutation (GSH), ascorbic acid (AA), uric acid (UA), and ceruloplasmine (CP). Glucose level was determined using kits supplied by (Randox, UK). Total cholesterol TC, Triglycerides were determined using kits from (spinreact, Spain). Uric acid was measured by enzymatic colorimetric assay using kits supplied by (Randox, UK). Hemoglobin A<sub>1c</sub> program intended for the determination of Glycated hemoglobin (A<sub>1c</sub>) in human depended on high performance liquid chromatography and who supplied by Variant Company, USA. Plasma malondialdehyde was measured according to the method described by Shah and Walker<sup>[16]</sup>. using anticoagulant tubes provided with EDTA. Estimation of erythrocyte glutathione was based on the method described by Butler *et al*<sup>[17]</sup>. using anticoagulant tubes provided with acid – citrate-dextrose (ACD). Measurement of whole blood ascorbic acid was done according to the method described by Tetiz *et al*,<sup>[18]</sup> using anticoagulant tubes provided with heparin. Serum ceruloplasmin was measured according to Mancini method<sup>[19]</sup> by using single radial immune diffusion (RID) endplates; supplied by Biomegreb, Tunisia.

**Statistical analysis:**

Data were analyzed using computer facility-the available statistical packages of SPSS-11 (statistical packages for social sciences-version 11.0). Data were present in simple measures of Mean±SD. The significance of difference between quantitative variables was tested using student t-test for comparing between two means of independent groups. P value equal and less than 0.05 was used as the level of significance, and P value equal and less than 0.01 was used as the level of a highly significant. Pearson correlation coefficient is significant at the 0.05 level (2-tailed)

**Results:**

The current study showed that there was a highly significant increased (P<0.001) in fasting serum glucose FSG, glycated hemoglobin (HbA<sub>1c</sub>), and malondialdehyde (MDA) in poorly controlled diabetic patients group G2 comparing to well controlled diabetic patients G1(189.91±22.58 vs 103.3±16.23; 9.1 ± 1.2 vs 6.01± 0.9; 8.9±1.1 vs 6.8±0.6) respectively; although,

there was a significant increased increase (P<0.05) in total cholesterol (TC), and triglycerides (181.88±6.19 vs 164.62±5.2; 198.36±9.07 vs 160.31 ±4.1) respectively. Instead of that; there was significant increase in Ceruloplasmine levels (CP) (P<0.05) in poorly controlled patients. On the other hand, there was a highly significant decreased in (erythrocyte-GSH, ascorbic acid (AA) in poorly controlled diabetic patients compared to well controlled diabetic patients (41.7±5.04 vs 67.18±10.13, and 1.11±0.1 vs 1.71±0.2) correspondingly; and there was significant decreased (P<0.05) in uric acid (UA)(3.63±1.05 vs 5.05±1.1). Correlation analysis showed a highly significant positive correlation in between HbA<sub>1c</sub> and FSG (P<0.01) only in well controlled patients group, and significant positive correlation (P<0.05) between HbA<sub>1c</sub> and TC only in poorly controlled group. As well as, there is a significant positive correlation between HbA<sub>1c</sub> with MDA, and erythrocyte-GSH with AA in both group. On the other hand, there was a significant negative correlation in MDA-GSH, MDA-AA, MDA-UA in both studying group.

**Table-1: Clinical characteristics and parameters of type 2 DM according to glycemic control and healthy subjects.**

| Clinical characteristics | Type 2 DM   |              | P value   |
|--------------------------|-------------|--------------|-----------|
|                          | G1          | G2           |           |
| No.                      | 75          | 70           | -         |
| Age (yrs)                | 47.6±10.3   | 48.01± 12.4  | N.S       |
| Sex (m:f)                | (36: 37)    | (35: 38)     | N.S       |
| BMI (kg/m <sup>2</sup> ) | 26.95±5.6   | 27.33±3.5    | N.S       |
| Duration (yrs)           | 7.98±0.8    | 10.1± 1.2    | <0.05*    |
| FSG (mg/dl)              | 103.3±16.23 | 189.91±22.58 | <0.0001** |
| HbA <sub>1c</sub>        | 6.01± 0.9   | 9.1 ± 1.2    | <0.001**  |
| MDA (µmol/L)             | 6.8±0.6     | 8.9±1.1      | <0.001**  |
| Erythrocyte-GSH (µmol/L) | 67.18±10.13 | 41.7±5.04    | <0.0001** |
| AA (mg/dl)               | 1.71±0.2    | 1.11±0.1     | <0.01**   |
| UA (mg/dl)               | 5.05±1.1    | 3.63±1.05    | <0.05*    |
| CP (mg/dl)               | 43.4±7.47   | 51.4±9.59    | <0.05*    |
| TC (mg/dl)               | 164.62±5.2  | 181.88±6.19  | <0.05*    |
| TG (mg/dl)               | 160.31 ±4.1 | 198.36±9.07  | <0.05*    |

Data are mean ± SD, \*P ≤ 0.05 was considered significant, \*\*P < 0.01 is a highly significant. N.S non significant.

Table-2: Correlation between parameters in type 2 DM according to glycemic control.

| Parameters | G1      | G2      |
|------------|---------|---------|
|            | r       | r       |
| HbA1c-FSG  | 0.62**  | 0.16    |
| HbA1c-TC   | 0.19    | 0.44*   |
| HbA1c-TG   | 0.15    | 0.23    |
| HbA1c-MDA  | 0.42*   | 0.52**  |
| MDA-GSH    | -0.71** | -0.77** |
| MDA-AA     | -0.78** | -0.69** |
| MDA-UA     | -0.40*  | -0.44*  |
| GSH-AA     | 0.47*   | 0.51*   |

\*correlation is significant at the 0.05 level (2-tailed)

\*\*correlation is a highly significant at the 0.01 level (2-tailed)

### Discussion:

Diabetes mellitus has been known to be a state of excess generation of free radicals which are contributed by several mechanisms, including hyperglycaemia and a defective antioxidant status, which causes oxidative stress<sup>[11]</sup>. The data of the present study revealed increased levels of lipid peroxidation marker described as (MDA) and a depleted level of the antioxidant status in poorly controlled type 2 diabetics, in favour of an oxidative stress in such patients. These results were in agreement with those of previous studies<sup>[5,11]</sup>, which demonstrated a strong association between poor glycaemic control and the depletion of the protective antioxidant defense mechanisms in diabetes mellitus. But others<sup>[20, 21]</sup> failed to observe lipid peroxidation association with glycemic control. This discrepancy may result from the differences in disease duration and / or the age of subjects under evaluation, since the levels of HbA1c have been age dependent and disease duration related<sup>[22]</sup>. In the present study, all the diabetic patients who are poorly controlled observed hyperlipidaemia, which

was in agreement with the findings of other studies<sup>[23,25]</sup>. These findings are not surprising, because long-term hyperglycaemia causes generalized vascular endothelial damage, which reduces functional lipoprotein lipase, leading to an increase in TC, TG and a decrease in HDL-C.<sup>[25]</sup> Several studies suggested that improvement in glycemic control might reduce both microvascular and macrovascular complications of diabetes<sup>[26,27]</sup>. Other reports postulated a reduction in oxidative stress achieved by optimal glycemic control<sup>[28,29]</sup>. Malondialdehyde is one of the lipid peroxidation products frequently used to determine the oxidant/antioxidant balance in diabetic patients<sup>[30,31]</sup>. In agreement with other finding<sup>[32]</sup>, the present study demonstrated that well glycemic control patients had lower MDA level compared to poor-glycemic control group, which may confirm the role of glycemic control to decrease levels of lipid peroxidation by-products, even when these levels were not normalized. It was suggested that hyperglycemia enhance the generation of free radicals through several biochemical pathways (non-enzymatic glycation, polyol

pathway, and glucose autoxidation)<sup>[29]</sup>. Free radicals can result in consumption of antioxidant defenses and increase susceptibility to lipid peroxidation<sup>[29]</sup>. Hyperglycemia in diabetes can generate free radicals, and hydrogen peroxide, which cause membrane lipid peroxidation and osmotic fragility in human red blood cells (RBCs). Mechanisms that have been proposed to explain the glutathione effect include the removal of species that initiate lipid peroxidation, scavenging of radicals, scavenging of peroxy radicals and maintenance of membrane protein thiols<sup>[33]</sup>. Thus GSH may be exhausted through FRs scavenging and forming of oxidized glutathione (GSSG), which is consistently being reduced by glutathione disulfide reductase (GSSG reductase). Consistent with other findings<sup>[34]</sup> good glycemic control had significantly higher level of Ery-GSH compared to poor glycemic control, which support the hypothesis that glycemic control has a potential role to restore GSH depletion. Although, Hatice *etal*<sup>[31]</sup>. explain that hyper glycaemic conditions, glucose is preferentially used in the polyol pathway that consumes NADPH which is necessary for GSH regeneration by the glutathione reductase enzyme. Hyperglycemia is therefore indirectly the cause of GSH depletion. As GSH is an important molecule, its depletion leads to the increase of oxidative stress. Another finding in the present study showed that well-glycemic control had significant higher level of AA compared to other diabetic group, which support the hypothesis to the effect of glycemic control to improve antioxidant defense systems<sup>[26]</sup>. This results Consistent

with other<sup>[25, 35]</sup>. In addition to that, the results documented the hypothesis that glycemic control could restore depletion of antioxidants in diabetic patients<sup>[36]</sup>. Firoozrai *etal*<sup>[37]</sup> clarify that ascorbic acid metabolism is altered in conditions such as hyperglycemia and insulin insufficiency and patients with diabetes have about 30% lower plasma ascorbic acid levels than persons without diabetes. Decrease in active transport of reduced ascorbic acid and uptake inhibition of the oxidized form has observed in hyperglycemia<sup>[37]</sup>. Uric acid considered one of antioxidants that directly scavenging oxygen radicals, singlet oxygen, oxo-haem oxidants and hydroxyl radicals, another important antioxidant property of uric acid is its ability to inhibit ascorbate oxidation, as well as lipid peroxidation. In contrast to other antioxidant scavenger reactions, the inhibition of ascorbate oxidation and lipid peroxidation provided by uric acid does not involve uric acid oxidation<sup>[38]</sup>. Obviously G1 had higher level of uric acid compared to G2, which is in somehow related to well glycemic control of group1 compared to the poor glycemic control of G2, no available study for the effect of glycemic control on uric acid, but several studies investigate the antioxidants status including uric acid in uncomplicated diabetics which revealed a significant reduction in uric acid levels patients with short duration of disease and without diabetic complications<sup>[39]</sup>. Cerulo plasmin (CP) is one of plasma proteins that acts as a potent antioxidant, with different physiological roles include scavenge super oxide anion radicals ( $O_2^{\cdot-}$ ). Through this ability CP acts as a major extracellular

scavenger<sup>[40]</sup>. CP levels in our study is linked somehow to the degree of glycemic control, as glycemic state worsen and lead to increase CP levels in diabetic patients. The correlations in this study showed a positive correlation between A1c and FSG demonstrated in diabetic G1, was helpful to understand that fasting glycemia provide evidence for the metabolic control achieved by HbA1c<sup>[41]</sup>. Lack of such correlation in G2 may be due to the disturbance in glycemia state of this group. Although, it is important to recognize the associations between parameters as these do not function in isolation<sup>[42]</sup>. In the current study poor glycemic control was associated with enhanced oxidative stress as indicated by a strong correlation between glycated hemoglobin and marker of lipid peroxidation (MDA) in diabetic patients<sup>[5,42]</sup>. The positive correlations between Ery-GSH and AA were found in both study groups, this was supported by previous studies<sup>[27-29]</sup>. These studies have suggested that in diabetes GSH is largely consumed, mainly because of the regeneration of AA, which is extensively oxidized in diabetic patients. Thus existence of such correlation in both study groups provided useful evidence for the measurement of AA in whole blood to be a better index than plasma. MDA presented in this study as lipid peroxidation by-product was negatively correlated with AA<sup>[26]</sup>, Ery-GSH<sup>[28,29]</sup>, and uric acid in both groups. The present study concluded that these correlations may reflect a significant increase in lipid peroxidation (MDA) accompanied by a significant decrease in the antioxidant mechanism in poorly controlled group by the inverse relationship between them. As a

result, glycemic control has a potential role to lower lipid peroxidation by-products, and reduce consumption of antioxidants. It may be appropriate to evaluate markers of oxidative stress in addition to routine laboratory assessments in evaluation of type2 DM patients. Antioxidant supplementation may be necessary for treatment to reduce oxidative stress for diabetic complications protection and amelioration.

### **References:**

- 1- Stratton, I. M.; Adler, A. I.; Neil, H. A.; Matthews, D. R.; Manley, S. E. Cull, C. A. Hadden, D.; Turner, R. C. and Holman, RR. Association of glycemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ*. 2000. Vol. 321 (7258). Pp: 405-12.
- 2- Brownlee, M. Biochemistry and molecular cell biology of diabetic complications. *Nature*. 2001. Vol. 414 (6865). Pp: 813-20.
- 3- The Diabetes Control and Complications Trial Research Group. The relationship of glycemic exposure (HbA1c) to the risk of development and progression of retinopathy in the diabetes control and complications trial. *Diabetes*. 1995. Vol. 44 (8). Pp: 968-83.
- 4- Hirsch, I. B. and Brownlee, M. Should minimal blood glucose variability become the gold standard of glycemic control? *J. Diabetes Complications*. 2005. Vol. 19 (3). Pp: 178-81.
- 5- Shadia, A. F.; Safinaz, A.; Wafaa, I. R.; Azza, S. H.; Gamila, S. M.; Omima, M. S. And Safaa, M. Y. Diagnostic Potential of oxidative Stress Indicators in Egyptian Type 1 Diabetic Children and Adolescents. *Journal of Applied Sciences Research*. 2009. Vol. 5 (12). Pp: 2524-2534.

- 6- Sarah, E. S.; Temo, B.; Wim, K.; Joost, B. Hoekstra, M. and Hans, D. V. No Relevant Relationship between Glucose Variability and Oxidative Stress in Well-Regulated Type 2 Diabetes Patients. *Journal of Diabetes Science and Technology*. 2011. Vol. 5 (1). Pp: 86-92.
- 7- Pasupathi, P. V.; Chandrasekar and Kumar. S. U.; Evaluation of oxidative stress, enzymatic and non-enzymatic antioxidants and metabolic thyroid hormone status in patients with diabetes mellitus. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews*. 2009. Vol.3. Pp: 160-165.
- 8- Ramakrishna, V.; and Jailkhani, R.; Evaluation of oxidative stress in IDDM patients. *Diagnostic Pathology*. 2007. Vol. 2. Pp: 22.
- 9- Ferdinando, G.; and Michael, B.; Oxidative Stress and Diabetic Complications. *Circ Res*. 2010. Vol. 107. Pp: 1058-70.
- 10- Krzystek-Korpaczka, M.; Salmonowicz, B.; Boehm, D.; Berdowska, I.; Zielinski, B.; Patryn, E.; Noczynska, A. and Gamian, A. Diagnostic potential of oxidative stress markers in children and adolescents with type 1 diabetes. *Clinical Biochemistry*. 2008. Vol. 41. Pp: 48-55.
- 11- Bonnefont-Rousselot, D.; Bastard, J. P.; Jaudon, M. C. and Delattre, J. Consequences of the diabetic status on the oxidant/antioxidant balance. *Diab & Metbol*. 2000. Vol. 26. Pp: 163-76.
- 12- Maritim, A. C.; Sanders R. A. and Watkins JB. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol*. 2003. Vol. 17. Pp: 24-38.
- 13- Vadde, R. and Rama, J.; Evaluation of oxidative stress in insulin dependent diabetes mellitus (IDDM) patients. *Diagnostic Pathology*. 2007. Vol. 2. Pp: 22
- 14- Martin-Gallan, P.; Carrascosa, A.; Gussinye, M.; and Dominguez, C. Biomarkers of diabetes associated oxidative stress and antioxidant status in young diabetic patients with or without subclinical complications. *Free Radical Biology and Medicine*. 2003. Vol. 34. Pp: 1563-74.
- 15- Zwart, L.; Meerman, N.; Commadeur, M. and Vermeulen, E. Biomarkers of free radical damage ,applications in experimental animals and in humans. *Free Radical Biology and Medicine*. 1999. Vol. 26 (1/2). Pp: 202-226.
- 16- Shah S.V.; Walker, P.D. and Evidence suggesting a role for hydroxyl radical in glycerol induced acute renal failure. *Am-J-Physiol*. 225, ( Renal,fluid, electrolyte physiol 24) 1988. Pp: 438-443.
- 17- Butler E.; Duron D. and Kelly B. M.; Improved method for the determination of blood glutathione. *J-Lab-Clin-Med*. 1963. Vol. 6. Pp: 882-888.
- 18- Tetiz, N. w. Textbook of clinical chemistry. W. B. Saunders Company. Philadelphia.
- 19- Mancini, G.; Carbouara A. O. and Heremous J. F. Immunochemical quantitation of antigens by signal radial immune diffusion. *Immuno chemistry*. 1965. Vol. 2. Pp: 235-254.
- 20- Martin- Gallan, P.; Carrascosa, A.; Gussinye, M.; and Dominguez, C.; Biom arkers of diabetesassociated oxidative stress and antioxidant status in young diabetic patients with or without subclinical complications. *Free Radical Biology and Medicine*. 2003. Vol. 34. Pp: 1563-74.
- 21- Füsün, E.; Fatma, T.; Banu, A.; and Yasemin, U. Glycemic control, oxidative stress, and lipid profile in children with type 1 diabetes mellitus. *Archives of Medical Research*. 2004. Vol. 35 (2). Pp: 134-140.



- 22- Levine, B. S.; Anderson, B. J.; Butler, D. A.; Antisdel, J. E.; Brackett, J. and Laffel, LM. Predictors of glycemic control and short-term adverse outcomes in youth with type 1 diabetes. *Journal of Pediatric*. 2001. Vol. 139. Pp: 197-203.
- 23- Suryawanshi, N. P.; Bhutey, A. K.; Nagdeote, A. N.; Jadhav, A. A., and Manoorkar, GS. Study of lipid peroxide and the lipid profile in diabetes mellitus. *Indian J of Clin Biochem*. 2006. Vol. 21. Pp: 126-30.
- 24- Goldberg, J. Clinical review 124: Diabetic dyslipidemia: causes and consequences. *J Clin Endocrinol Metab*. 2001. Vol. 86. Pp: 965-71.
- 25- Suchitra, M. M; Pallavi, M.; Shivaprasad, P.; Sachan, A.; Madhusudhana, R. A, Aparna, R. B. and Srinivasa, Rao. Uric Acid, Oxidants and Antioxidant Vitamins in Type 2 Diabetic Patients without Cardiovascular Complications. *J. Clin and Diagnostic Res*. 2011. Vol. 5 (6). Pp: 1161-4.
- 26- Ceriello, A. Oxidative Stress and glyce mic control. *Metabolism*. 2000. Vol. 49 (2.1). Pp: 27-29.
- 27- Seghieri, G.; Disimplicio, P.; DeGiorgio L. A.; Aanichini R. and Franconi F.. Relationship between metabolic control and platelet content of glutathione and its related enzymes, in insulin-dependent diabetes mellitus. *Clin-Chem-Acta*. 2000. Vol. 299 (1-2). Pp: 109-117.
- 28- Sharma A.; Kharb S.; Chugh S.N.; Kakkar R. and Singh, G.P. Effect of glycemic control and vitamin E supplementation on total glutathione content in non-insulin-dependent diabetes mellitus. *Ann.Nutr.Metab*. 2000. Vol. 44 (1). Pp: 11-13.
- 29- Sharma, A.; Kharb, S.; Chugh S. N.; Kakkar, R.; and Singh, G.P. Evaluation of oxidative stress before and after control of glycemia and after vitamin E supplementation in diabetic patients. *Metabolism*. 2000. Vol. 49 (2). Pp: 160-162.
- 30- Altomare, E.; Vendemiale, G.; Chicco, D.; Procacci, V. and Cirelli, F. Increased lipid peroxidation in type 2 poorly controlled diabetic patients. *Diabetes Metab*. 1992. Vol.18. Pp: 264-271.
- 31- Hatice, P.; Banu, S.; and Neslihan, B. Lipid Peroxidation and Resistance to Oxidation in Patients with Type 2 Diabetes Mellitus. *Tohoku J. Exp. Med*. 2004. Vol. 203. Pp: 211-18.
- 32- El-ghoroury, E. A.; Raslan, H. M.; Badawy, E. A.; El-Saaid, G. S.; Agybi, M. H.; Siam, I. and Salem, SI. Malondialdehyde and coen zyme Q10 in platelets and serum in type 2 diabetes mellitus: correlation with glycemic control. *Blood Coagulation & Fibrinolysis*. 2009. Vol. 20 (4). Pp: 248-251.
- 33- Jain, S. K.; McVie, R. and Smith, T.; Vitamin E supplementation restores glutathione and malondialdehyde to normal concentrations in erythrocytes of type 1 diabetic children. *Diabetes Care*. 2000. Vol. 23 (9). Pp: 1389-1394.
- 34- Muruganandam, A.; Drouillard, C.; Thibert, R. J.; Cheung R. M.; Draisey, TF. and Mutus, B. Glutathione metabolic enzyme activities in diabetic platelets as a function of glycemic control. *Thromb Res*. 1992. Vol. 67. Pp: 385-397.
- 35- Ganesh, N. D.; Harshal, V. C.; and Meena, S. Supplementation of Vitamin C Reduces Blood Glucose and Improves Glycosylated Hemoglobin in Type 2 Diabetes Mellitus: A Randomized, Double- Blind Study. *Adv Pharmacol Sci*. 2011. Pp: 195-271.

## AJPS, 2013, Vol. 14, No.2

- 36- Dosoo, D. K.; Rana, S. V.; Offe-Amoyawk. Tete-Donkor, D. Maddy SQ. Total antioxidants status in type 2 diabetic patients in Ghana. *Diabetes international*. 2000. Vol.10 (1). Pp: 26-27.
- 37- Firoozrai, M.; Nourmohammadi, I. and Khanaki, K. Assessment of Antioxidant Vitamins Retinol and  $\alpha$ -Tocopherol in Plasma and Ascorbic Acid in Plasma and Mononuclear Leukocytes in Type 2 Diabetics. *Int J. Endocrinol Metab*. 2006. Vol.4. Pp: 202-205
- 38- Davies, K. J.; Seranian, A.; Muakkassah K. F. S.; Hochstein P. Uric acid-iron ion complexes. A new aspect of the antioxidant function of uric acid. *Biochemical Journal*. 1986. Vol. 235. Pp: 747-754.
- 39- Marra, G.; Cotroneo, P.; and Pitocco, D. *et al.* Early increase of oxidative stress and reduced antioxidant defenses in patients with uncomplicated type 1 diabetes. *Diabetes Care*. 2000. Vol. 25 (2). Pp: 370-375.
- 40- Telci, A.; Cakatay, U.; Kayali, R.; and Erdogan, C. *et al.* Oxidative protein damage in plasma of type 2 diabetic patients. *Horm- Metab- res*. 2000. Vol. 32 (1). Pp: 40-43.
- 41- Ko, G. T. C.; Chan, J.C.N.; Tsang, L.W.W; Cockram C.S. Combined use of fasting plasma glucose and HbA1c predicts the progression to diabetes in Chinese subjects. *Diabetes Care*. 2000. Vol. 23 (12). Pp: 1770-1773.
- 42- Matteucci, E. and Giampietro, O.; Oxidative stress in families of type 1 diabetic patients. *Diabetes Care*. 2000. Vol. 23 (8). Pp: 1182-1186.